

Alpha-decay Rates of Yb and Gd in Solar Neutrino Detectors

M. Fujiwara,^{1,2} T. Kawabata,³ P. Mohr⁴

¹ *Research Center for Nuclear Physics, Osaka University, Mihogaoka 10-1, Ibaraki, Osaka
567-0047, Japan*

² *Advanced Science Research Center, JAERI, Tokai, Ibaraki, 319-1195 Japan*

³ *Department of Physics, Kyoto University, Kyoto 606-8502, Japan*

⁴ *Institut für Kernphysik, Technische Universität Darmstadt, Schlossgartenstrasse 9, D-64289
Darmstadt, Germany*

(February 8, 2008)

Abstract

The α -decay rates for the nuclides $^{168,170,171,172,173,174,176}\text{Yb}$ and $^{148,150,152,154}\text{Gd}$ have been estimated from transmission probabilities in a systematic α -nucleus potential and from an improved fit to α -decay rates in the rare-earth mass region. Whereas α -decay of ^{152}Gd in natural gadolinium is a severe obstacle for the use of gadolinium as a low-energy solar-neutrino detector, we show that α -decay does not contribute significantly to the background in a ytterbium detector. An extremely long α -decay lifetime of ^{168}Yb is obtained from calculation, which may be close to the sensitivity limit in a low-background solar neutrino detector.

PACS numbers: 23.60.+e, 26.65.+t, 27.60.+j, 27.70.+q

The solar neutrino problem is an important current subject to be studied in relation to the fundamental physics of neutrino oscillations. After the successful measurements of the solar neutrinos by Super-Kamiokande [1], SAGE [2], GALLEX [3], and SNO [4], the problem appears to be solved best in terms of oscillation of ν_e into other neutrino flavors. To arrive at the final solution of the solar- ν problem, a real-time measurement of the ν_e -energy spectrum including pp -, ${}^7\text{Be}$ -neutrinos is now of central interest [5].

The solar neutrino detection requires an extremely low-background measurement with new technologies. Raghavan [6] has suggested a flavor-specific detection scheme with low thresholds for the real-time detection of solar neutrinos. The neutrino captures $\nu_e + {}^{176}\text{Yb} \rightarrow e^- + {}^{176}\text{Lu}^*$ and $\nu_e + {}^{160}\text{Gd} \rightarrow e^- + {}^{160}\text{Tb}^*$ are based on charged current mediated Gamow-Teller transitions. For the identification of a neutrino capture event a delayed coincidence between a prompt e^- and a γ -ray from the decay of a nanosecond isomer in ${}^{176}\text{Lu}$ or ${}^{160}\text{Tb}$ is used. For both suggested nuclei, ${}^{176}\text{Yb}$ and ${}^{160}\text{Gd}$, one finds sufficient Gamow-Teller strength at low energies and a nanosecond isomer which is populated in the neutrino capture reaction. Typical detection rates of solar neutrinos are of the order of ten counts per year and per ton of detector material [7].

A new solar neutrino detector using Yb materials has been proposed in LGNS at Gran Sasso [6–8], and sources of background for such a detector are presently being studied [9]. The α -decay from the material in a solar- ν detector is a possible serious problem to pollute the spectrum consisting of rare neutrino events. For example, it has turned out that Gd material is not suitable for a solar- ν detector; the ${}^{152}\text{Gd}$ isotope decays by α emission with a half-life of $T_{1/2} = 1.08 \times 10^{14}$ years, and is followed by the subsequent ${}^{148}\text{Sm}$ α -decay with $T_{1/2} = 7 \times 10^{15}$ years. The number of ${}^{152}\text{Gd}$ atoms included in the natural Gd metal with 20 tons amounts to 1.5×10^{26} atoms. The decay rate calculated by the well-known radioactive decay formula

$$\frac{dN}{dt} = -N_0 \lambda e^{-\lambda t} \quad (1)$$

amounts to about 30,000 counts/sec for the 20 ton Gd material. This is a huge decay rate,

which decreases the reliability of the coincidence events due to a large number of background α -signals and due to accidental coincidences even in the case of a narrow time gate less than 10^{-8} seconds for coincidence.

Recently, ytterbium has been suggested as material for a solar- ν detector. The Q-values of α -particle emission for all the Yb isotopes are positive, but relatively small. Hence, each Yb isotope must have an extremely long half-life although there has been no report on the α -decay lifetime in literature. Such half-lives may be measurable with a large mass detector like a solar- ν detector. If the half-life of some Yb isotope is relatively short, the α -decay may cause a significant background in the analysis of the obtained neutrino data. Therefore, it is important to estimate the α -decay rates of the Yb isotopes and to compare them with the solar neutrino rates. Recently, Gamow-Teller strengths in the inverse β transition $^{176}\text{Yb} \rightarrow ^{176}\text{Lu}$ were studied, and it has been concluded that Yb-based detectors are well-suited for the real-time spectroscopy of the sub-MeV solar neutrinos [7,10].

There are several parameterizations for α -decay rates available mainly for nuclei in the region of $Z \geq 84$ [11–13]. The predictions from these parameterizations usually show small deviations for experimentally available decay rates. However, there often happen large discrepancies, up to ten orders of magnitude, between the predictions for extremely small decay rates (corresponding to small Q_α values). In this report, we wish to discuss the α -decay rates of Yb and Gd isotopes in the rare earth nuclei.

The lifetime of α -decay was first formulated by Geiger and Nuttall [14]. The probability of α -particle emission, W , can be expressed as

$$W = p_\alpha \nu T, \quad (2)$$

where p_α is the probability of finding an α -particle in a nucleus, ν the frequency of an α -particle appearing at the wall of a nucleus, and T is the transmission coefficient for the tunnelling effect. The half-life $T_{1/2}$ can be parameterized [15] as

$$\log_{10} T_{1/2} = a_1 + \log_{10} \frac{\sqrt{Q_\alpha}}{A^{1/3}} + a_2 \sqrt{Z_d A^{1/3}} + a_3 \frac{Z_d}{\sqrt{Q_\alpha}} \quad , \quad (3)$$

with the half-life $T_{1/2}$ in years, the Q -value for α -decay Q_α in MeV, and the mass (proton, neutron) numbers A (Z, N) of the parent nucleus and $A_d = A - 4$ ($Z_d = Z - 2, N_d = N - 2$) of the daughter nucleus.

Since the data fittings in most of Refs. [11–13] are focused on the α -decay for the nuclei with $Z > 80$, we tried to fit the α -decay half-lives for nuclei in the rare earth region separately. The data for the half-lives of the α -decay nuclei in the rare earth region were fitted using Eq. (3) with a_1 , a_2 , and a_3 as three parameters. All Q_α values have been taken from the mass table of Audi and Wapstra [16]. The best fitting parameters, a_1 , a_2 , and a_3 are $a_1 = -34.577$, $a_2 = -1.2685$, and $a_3 = 1.7336$, respectively. The α -decay half-lives are found to be well fitted as can be seen from Fig. 1. The χ^2 per degree of freedom has been significantly improved. In our 3-parameter fit χ^2 is reduced by a factor of about 4 compared to the 2-parameter fit of Ref. [11], and by a factor of about 2 compared to the 2-parameter fit of Refs. [12,13] which is based on the model of Bethe [17]. These 2-parameter fits are given by [11]

$$\log_{10}T_{1/2} = (9.54Z_d^{0.6}/\sqrt{Q_\alpha}) - 58.87 \quad (4)$$

and [12,13]

$$\log_{10}T_{1/2} = 1.598(Z_d/\sqrt{Q_\alpha} - Z_d^{2/3}) - 27.44 \quad (5)$$

Tables I and II show the expected half-lives for the Yb and Gd isotopes. The experimental half-life of 1.1×10^{14} years for ^{152}Gd agrees with the result of the present fitting within a factor of 1.4. However, the predictions from the two parameter fittings by Brown [11] and Refs. [12,13] differ by three orders of magnitude. A similar agreement is also found for ^{148}Gd and ^{150}Gd . Again there are strong deviations between the predictions mainly given from the fits of α -decay rates for nuclei with $Z \geq 84$ [11–13]. Here, one can address the question that the discrepancy may come from the deformation effect on α -decay for the rare earth nuclei. But, it should be noted that the most nuclei with $Z > 84$ are deformed as well, and the influence of nuclear deformation does not change the half-life by an order of magnitude (see discussion below).

Our prediction for the different quasi-stable Yb isotopes varies from $T_{1/2} = 4.5 \times 10^{24}$ years for ^{168}Yb to $T_{1/2} = 6.2 \times 10^{95}$ years for ^{176}Yb . The parameterizations from literatures are between one order of magnitude (^{168}Yb) smaller [12,13] (larger [11]) to five orders of magnitude (^{176}Yb) smaller [12,13] (larger [11]). Note that no useful prediction for the relevant α -decay half-life of ^{168}Yb can be derived from Refs. [11–13] because their predictions differ by more than three orders of magnitude. Of course, the calculated half-lives of $^{172,173,174,176}\text{Yb}$ exceed any measurable range. But, the excellent agreement between the new fit and the half-lives from the folding potential calculations (see below) gives us some confidence that our predictions are reliable in a very broad range of half-lives (where other extrapolations show huge deviations).

Because of these large discrepancies between the extrapolations of the various parameterizations, we applied the semi-classical model of Gurvitz and Kälbermann [18] together with the systematic folding potentials of Atzrott *et al.* [19]. The required nuclear densities were derived from measured charge density distributions [20]. When no measured charge distribution was available, the measured distribution of a neighboring isotope was used with an adjusted radius parameter $R \sim A^{1/3}$. This model was already successfully used to calculate the α -decay properties of neutron-deficient so-called *p*-nuclei in [21].

In the semi-classical model, the α -decay width Γ_α is given as [18]:

$$\Gamma_\alpha = PF \frac{\hbar^2}{4\mu} \exp \left[-2 \int_{r_2}^{r_3} k(r) dr \right] \quad (6)$$

with the preformation factor P , the normalization factor F

$$F \int_{r_1}^{r_2} \frac{dr}{k(r)} = 1 \quad (7)$$

and the wave number $k(r)$

$$k(r) = \sqrt{\frac{2\mu}{\hbar^2} |Q_\alpha - V(r)|} \quad , \quad (8)$$

where μ , Q_α , and r_i are the reduced mass, the α -decay energy, and the classical turning points, respectively. For $0^+ \rightarrow 0^+$ *s*-wave decay, the inner turning point is at $r_1 = 0$. r_2

varies from about 7 to 9 fm, and r_3 varies up to about 350 fm for the lowest Q_α value. The decay width Γ_α is related to the half-life by the well-known relation $\Gamma_\alpha = \hbar \ln 2 / T_{1/2}$. For the odd Yb isotopes, an additional centrifugal potential was used. The preformation factors P were estimated from the systematic trend of P shown in Fig. 2 of Ref. [21] leading to $P = (4.0 \pm 0.8)\%$ for the Yb isotopes and $P = (10.4 \pm 2.5)\%$ for the Gd isotopes. Larger P values can be expected around the shell closures $N = 82$ and $N = 126$, whereas smaller values for P are found in between. The results of this model are listed in Tables I and II. They are in good agreement with our new fits in the rare earth region. Therefore, the newly determined α -decay formula can be used for predictions in the rare earth region with very limited uncertainties.

A systematic study of α -decay half-lives [22] using the same model [18] but specially shaped potentials is in rough agreement with our results for the Gd isotopes (8th line in Table II). Unfortunately, the quasi-stable Yb isotopes are not considered in Ref. [22].

There are several sources of uncertainties for the presented predictions for the half-lives of the Yb and Gd isotopes. Because of the roughly exponential dependence of the tunneling probability on the α decay energy Q_α , a factor $f(\Delta Q_\alpha)$ has been calculated by which the half-lives are shorted (enlarged) if Q_α is increased (decreased) by its uncertainty ΔQ_α . This factor $f(\Delta Q_\alpha)$ is given in Tables I and II. Typical uncertainties remain within 40 % with the exception of ^{176}Yb where the very low Q_α and its 4 keV uncertainty lead to an uncertainty of $f(\Delta Q_\alpha) = 3.61$. This uncertainty is common to all predictions in this work and also in Refs. [11–13].

The uncertainties of the predictions from the 3-parameter fit in this work are smaller than in the previous 2-parameter fits because of the improved χ^2 value. The average deviation for $\log_{10} T_{1/2}$ in our fit is about 0.5 corresponding to an average deviation of a factor of 3. Therefore, a typical uncertainty of a factor of 3 can be expected for our predicted half-lives. The uncertainties in the coefficients a_1 , a_2 , and a_3 lead to minor uncertainties compared to the average deviation of a factor of 3.

In the case of the folding potential calculations, the predicted values depend sensitively

on the average preformation factors derived from the systematics in Ref. [21]. Typical uncertainties of P are of the order of 20 – 30 % which lead to identical uncertainties for the predicted half-lives. Together with the uncertainties of Q_α , this leads to an overall uncertainty of better than a factor of two for the folding potential predictions.

It is well-known that nuclei in the rare earth region show strong deformations with typical deformation parameters in the order of $\beta_2 \approx 0.3$ [23] which may lead to additional uncertainties. The influence of deformation on the decay probabilities has been analyzed recently in Refs. [24–27]. In Ref. [24], it is stated that various calculations of the half-lives agree within a factor of 4 with the spherical result. In Ref. [25], it is suggested that the deformation effect enhances the decay probability by roughly a factor of 2, whereas in Ref. [26] the decay probability for deformed nuclei is reduced by a factor less than 2. In Ref. [27], the deformation effect leads only to small variations of the half-lives. The good quality of our fit to the decay data in the rare earth region confirms that the influence of deformation on the α -decay half-lives remains relatively small compared to the uncertainties that arise from the predictions of our and of previous fits [11–13].

In summary, we estimated the half-lives of α -decay for the Gd and Yb isotopes. Our predictions from the new fit in the rare-earth region and from the folding potential calculations agree within their uncertainties discussed above, whereas previous fits show large deviations especially for low Q_α values and long half-lives. The “short” half-life of ^{152}Gd of $T_{1/2} = 1.1 \times 10^{14}$ years leads to serious background problems in a neutrino detector. The estimated half-life for ^{168}Yb is between 4.5×10^{24} years and 1.1×10^{25} years. If this predicted half-life is correct, the LENS detector with a mass of 20 tons of natural Yb will detect about 6~14 counts/year for α -particle emission with $Q_\alpha = 1.951$ MeV from ^{168}Yb decay, being negligible compared with the solar- ν events. This low rate causes definitely no serious background problem. However, the longest α -decay radioactivity so far measured in the past may be found in a low-background detector designed for a real-time solar- ν spectroscopy.

REFERENCES

- [1] Y. Fukuda et al., Phys. Rev. Lett. **81**, 1158 (1998).
- [2] J.N. Abdurashitov et al., Phys. Rev. C **60**, 055801 (1999).
- [3] W. Hampel et al., Phys. Lett. B **447**, 127 (1999).
- [4] Q.R. Ahmad et al., Phys. Rev. Lett. **87**, 071301 (2001).
- [5] J.N. Bahcall, *hep-ex/0106086*.
- [6] R.S. Raghavan, Phys. Rev. Lett. **78**, 3618 (1997).
- [7] M. Fujiwara *et al.*, Phys. Rev. Lett. **85**, 4442 (2000).
- [8] LENS (Low Energy Neutrino Spectroscopy) collaboration, Proposal to Laboratori Nazionali del Gran Sasso (LNGS), 1999.
- [9] J.-P. Meyer, LENS Internal Report 2001-06-22.
- [10] M. Bhattacharya, C. D. Goodman, R. S. Raghavan, M. Palarczyk, A. García, J. Rapaport, I. J. van Heerden, and P. Zupranski, Phys. Rev. Lett. **85**, 4446 (2000).
- [11] B. A. Brown, Phys. Rev. C **46**, 811 (1992).
- [12] R. Taagepera and M. Nurmia, Ann. Acad. Sci. Fenn. Ser. **A VI**, 78 (1961).
- [13] K. A. Keller and H. Z. Munzel, Z. Phys. **255**, 419 (1972).
- [14] H. Geiger and J. M. Nuttall, Philos. Mag. **22**, 613 (1911).
- [15] S. S. M. Wong, Introductory Nuclear Physics (Printice Hall, New Jersey, 1990) pp.233.
- [16] G. Audi and H. Wapstra, Nucl. Phys. **A595**, 409 (1995).
- [17] H. A. Bethe, Rev. Mod. Phys. **9**, 161 (1937).
- [18] S. A. Gurvitz and G. Kälbermann, Phys. Rev. Lett. **59**, 262 (1987).

- [19] U. Atzrott, P. Mohr, H. Abele, C. Hillenmayer, and G. Staudt, Phys. Rev. C **53**, 1336 (1996).
- [20] H. de Vries, C. W. de Jager, and C. de Vries, At. Data Nucl. Data Tables **36**, 495 (1987).
- [21] P. Mohr, Phys. Rev. C **61**, 045802 (2000).
- [22] B. Buck, A. C. Merchant, and S. M. Perez, At. Data Nucl. Data Tables **54**, 53 (1993).
- [23] S. Raman, C. H. Malarkey, W. T. Milner, C. W. Nestor jr., and P. H. Stelson, At. Data Nucl. Data Tables **36**, 1 (1987).
- [24] T. L. Stewart, M. W. Kermode, D. J. Beachey, N. Rowley, I. S. Grant, and A. T. Kruppa, Nucl. Phys. **A611**, 332 (1996).
- [25] D. S. Delion and R. J. Liotta, Phys. Rev. C **56**, 1782 (2000).
- [26] F. García, O. Rodríguez, M. Gonçalves, S. B. Duarte, O. A. P. Tavares, and F. Guzmán, J. Phys. G **26**, 755 (2000).
- [27] A. Dimarco, S. B. Duarte, O. A. P. Tavares, M. Gonçalves, F. García, O. Rodríguez, and F. Guzmán, Int. J. Mod. Phys. E **9**, 205 (2000).

FIGURES




FIG. 1. Ratio $R = T_{1/2}^{\text{calc}} / T_{1/2}^{\text{exp}}$ between the calculated half-lives $T_{1/2}^{\text{calc}}$ with Eq. (3) to the experimental half-lives $T_{1/2}^{\text{exp}}$ for nuclei in the rare earth region. The dotted lines connect isotopes from the same element. The error bars are only experimental. Additional errors from the uncertainty of Q_α are not included (see last lines of Tables I and II). The analyzed range of Q -values and half-lives is broad: Q -values cover the range from 1.9 MeV (^{144}Nd) to 7.0 MeV (^{175}Hg), and half-lives $T_{1/2}$ cover 26 orders of magnitude from milliseconds (^{158}W , ^{169}Pt) to more than 10^{15} years (^{144}Nd , ^{148}Sm , ^{174}Hf , ^{186}Os).

TABLES

TABLE I. Isotopic abundance (%) of Yb, α -decay energy (Q_α), the expected half-lives ($T_{1/2}$) in unit of year, and the uncertainty factor $f(\Delta Q_\alpha)$ from the Q value error. The parameters of Eq. (3) were adjusted to the half-lives of the nuclides shown in Fig. 1.

	^{168}Yb	^{170}Yb	^{171}Yb	^{172}Yb	^{173}Yb	^{174}Yb	^{176}Yb
(%)	0.13	3.05	14.3	21.9	16.1	31.8	12.7
Q_α (MeV)	1.95075	1.73764	1.55895	1.31030	0.94586	0.74009	0.57102
ΔQ_α (MeV)	0.00412	0.00146	0.00139	0.00139	0.00145	0.00160	0.00417
$T_{1/2}$ (exp.)	-	-	-	-	-	-	-
$T_{1/2}(\text{est.})^a$	$1.1 \cdot 10^{27}$	$1.4 \cdot 10^{32}$	$1.6 \cdot 10^{37}$	$8.5 \cdot 10^{45}$	$3.0 \cdot 10^{64}$	$3.8 \cdot 10^{80}$	$7.7 \cdot 10^{99}$
$T_{1/2}(\text{est.})^b$	$5.5 \cdot 10^{23}$	$2.4 \cdot 10^{28}$	$9.3 \cdot 10^{32}$	$7.4 \cdot 10^{40}$	$4.7 \cdot 10^{57}$	$1.8 \cdot 10^{72}$	$5.5 \cdot 10^{89}$
$T_{1/2}(\text{est.})^c$	$4.5 \cdot 10^{24}$	$4.1 \cdot 10^{29}$	$3.5 \cdot 10^{34}$	$1.1 \cdot 10^{43}$	$1.5 \cdot 10^{61}$	$8.4 \cdot 10^{76}$	$6.2 \cdot 10^{95}$
$T_{1/2}(\text{est.})^d$	$1.1 \cdot 10^{25}$	$6.9 \cdot 10^{29}$	$4.7 \cdot 10^{35}$	$1.0 \cdot 10^{43}$	$2.1 \cdot 10^{61}$	$1.9 \cdot 10^{76}$	$6.9 \cdot 10^{94}$
$f(\Delta Q_\alpha)$	1.24	1.09	1.10	1.13	1.23	1.40	3.61

^aParametrization of B. A. Brown [11].

^bParametrization of R. Taagepera and M. Nurmiä [12], K. K. Keller and H. Z. Munzel [13].

^cthis work, from fit to rare earth region.

^dthis work, from folding potential, using $P = (4.0 \pm 0.8) \%$.

TABLE II. Isotopic abundance (%) of Gd, α -decay energy (Q_α), the expected half-lives ($T_{1/2}$) in unit of year, and the uncertainty factor $f(\Delta Q_\alpha)$ from the Q value error. The parameters of Eq. (3) were adjusted to the half-lives of the nuclides shown in Fig. 1 including $^{148,150,152}\text{Gd}$.

	^{148}Gd	^{150}Gd	^{152}Gd	^{154}Gd
(%)	-	-	0.20	2.18
Q_α (MeV)	3.27121	2.80887	2.20458	0.91991
ΔQ_α (MeV)	0.00003	0.00630	0.00142	0.00115
$T_{1/2}$ (exp.)	$7.5 \cdot 10^1$	$1.8 \cdot 10^6$	$1.1 \cdot 10^{14}$	-
$T_{1/2}$ (est.) ^a	$7.6 \cdot 10^3$	$7.1 \cdot 10^8$	$3.7 \cdot 10^{17}$	$2.9 \cdot 10^{59}$
$T_{1/2}$ (est.) ^b	$2.0 \cdot 10^2$	$4.4 \cdot 10^6$	$1.8 \cdot 10^{14}$	$6.7 \cdot 10^{50}$
$T_{1/2}$ (est.) ^c	$2.6 \cdot 10^1$	$1.1 \cdot 10^6$	$1.5 \cdot 10^{14}$	$4.1 \cdot 10^{53}$
$T_{1/2}$ (est.) ^d	$6.0 \cdot 10^1$	$1.7 \cdot 10^6$	$1.4 \cdot 10^{14}$	$4.1 \cdot 10^{52}$
$T_{1/2}$ (est.) ^e	$2.6 \cdot 10^1$	$9.0 \cdot 10^5$	$6.7 \cdot 10^{13}$	-
$f(\Delta Q_\alpha)$	1.00	1.18	1.06	1.17

^aParameterization of B. A. Brown [11].

^bParameterization of R. Taagepera and M. Nurmiä [12], K. K. Keller and H. Z. Munzel [13].

^cthis work, from fit to rare earth region.

^dthis work, from folding potential, using $P = (10.4 \pm 2.5) \%$.

^efrom Ref. [22].